



Watkins, Adam J. and Sinclair, Kevin D. (2014) Paternal low protein diet affects adult offspring cardiovascular and metabolic function in mice. *American Journal of Physiology – Heart and Circulatory Physiology*, 306 (10). H1444-H1452. ISSN 1522-1539

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/38476/1/Watkins%20et%20al%202014%2C%20Am%20J%20Phys.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Paternal low protein diet affects adult offspring cardiovascular and metabolic function in mice.

Adam J. Watkins^{1,2} and Kevin D. Sinclair¹

5

¹School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough.
LE12 5RD, UK

²Aston Research Centre for Healthy Ageing. School of Life and Health Sciences. Aston University.
10 Birmingham. B4 7ET, UK

Abbreviated Running Title: Paternal low protein diet and adult offspring health in mice

15 **Corresponding Author:**

Adam J. Watkins

Aston Research Centre for Healthy Ageing. School of Life and Health Sciences. Aston University.
Birmingham. B4 7ET, UK

20 E mail: a.watkins1@aston.ac.uk

Tel: +44 (0)121 204 3700

25

Whilst the association between maternal periconceptional diet and adult offspring health is well characterised, our understanding of the impact of paternal nutrition at the time of conception on offspring phenotype remains poorly defined. Therefore, we determined the effect of a paternal preconception low protein diet (LPD) on adult offspring cardiovascular and metabolic health in mice. Male C57BL/6 mice were fed either normal protein diet (18% casein; NPD) or LPD (9% casein) for 7 weeks prior to mating. At birth, a reduced male:female ratio ($P=0.03$) and increased male offspring weight ($P=0.009$) were observed in litters from LPD compared to NPD stud males with no differences in mean litter size. LPD offspring were heavier than NPD offspring at 2 and 3 weeks of age ($P<0.02$). However, no subsequent differences in body weight were observed. Adult male offspring derived from LPD studs developed relative hypotension (decreased by 9.2 mmHg) and elevated heart rate ($P<0.05$), whilst both male and female offspring displayed vascular dysfunction and impaired glucose tolerance relative to NPD offspring. At cull (24 weeks), LPD males had elevated adiposity ($P=0.04$), reduced heart:body weight ratio ($P=0.04$) and elevated circulating TNF- α levels ($P=0.015$) when compared to NPD males. Transcript expression in offspring heart and liver tissue was reduced for genes involved in calcium signalling (*Adcy*, *Plcb*, *Prkcb*) and metabolism (*Fto*) in LPD offspring ($P<0.03$). These novel data reveal the impact of sub-optimal paternal nutrition on adult offspring cardiovascular and metabolic homeostasis, and provide some insight into the underlying regulatory mechanisms.

Keywords: Adult offspring health; cardiovascular dysfunction; developmental programming; metabolic homeostasis; paternal diet.

Introduction

Studies in humans and animal models have identified strong associations between adult
60 disease risk and environmental perturbations experienced during early development (21). Gamete
maturation and developmental events associated with fertilisation and pre-implantation embryo
development appear particularly sensitive to changes in environmental conditions (50). Studies
from a diverse range of model species including rat (24), sheep (40), and human populations (10)
have revealed similar phenotypic changes in offspring growth, cardiovascular and metabolic
65 homeostasis following maternal periconceptional environment manipulation. In the sheep, half
maintenance feeding prior to conception induced changes in blastocyst transcript levels for genes
associated with metabolic activity (33). In the sheep and cow, global maternal gestational
undernutrition has been shown to elevate fetal blood pressure (14) impair adult offspring glucose
tolerance (45) and affect offspring growth and adiposity in a sex specific manner (26). Similarly in
70 rodents, the feeding of a maternal low protein diet (LPD) during gestation induces significant
changes in offspring birth weight and growth (53), preferences for high-fat foods (6), insulin
resistance (16) and hypertension and vascular dysfunction (37).

Whilst our understanding of the developmental consequences of manipulating the maternal
75 environment is well defined, the impact of paternal physiology and nutritional status around
conception remains largely under-investigated. Studies in humans and mice have demonstrated that
increasing male BMI associates with reduced sperm motility (19), increased incidences of sperm
abnormality (23) and DNA fragmentation (11), and reduced pregnancy rates (17). In mice, offspring
metabolic profiles including hepatic lipid and cholesterol biosynthesis at weaning, serum glucose,
80 IGF-1 and corticosterone levels are altered in response to paternal LPD (9) or pre-mating fasting
(2). Consumption of a high fat (27) or high energy (36) diet impacts negatively on sperm motility,
DNA integrity and blastocyst developmental rates and impairs offspring pancreatic β -cell function
(28). In men, paternal obesity has been shown to associate with decreased blastocyst development

and live birth rate (5) and the DNA methylation status of the *IGF2* differentially methylated region
85 in the cord blood of newborn children (42).

Whilst these studies identify intergenerational transmission of metabolic disorders in young
offspring through sperm mediated mechanisms, the impact on adult offspring cardiovascular and
metabolic phenotype remains unknown. Therefore, the aim of our current study was to determine
90 the impact of a paternal LPD on well-defined markers of adult offspring cardiovascular and
metabolic health, focusing on analysis of adult offspring blood pressure, arterial function, in vivo
glucose tolerance and the expression of cardiovascular and metabolic regulatory genes.

Materials and Methods

95 *Animal Treatments*

All mice and experimental procedures were conducted using protocols approved by, and in
accordance with, the UK Home Office Animal (Scientific Procedures) Act 1986 and local ethics
committee at the University of Nottingham. Virgin male (9 week old) and female (5-9 week old)
C57BL/6 mice (Harlan Ltd, Belton, Leicestershire, UK) were maintained for 2 weeks at the
100 University of Nottingham's Bio Support Unit on a 07:00–19:00 light-dark cycle at a temperature of
20–22°C with *ad libitum* access to chow (2018 Teklad Global 18% Protein Rodent Diet; Harlan,
UK) and water. Weight matched male mice were housed singly and allocated to either a control
normal protein diet (NPD; 18 % casein, 42.5% maize starch, 21.3 % sucrose, 10% corn oil, 5%
cellulose; n = 8) or isocaloric (calories/gm) low protein diet (LPD; 9% casein, 48.5 maize starch,
105 24.3% sucrose, 10% corn oil, 5% cellulose; n = 8) offered *ad libitum* (Special Dietary Services Ltd,
UK; composition published previously (24, 52)) for 7 weeks prior to initiation of mating, and
maintained on respective diets until cull at 32 weeks of age (Figure 1).

Virgin, chow fed 7-9 week old C57BL/6 females were caged singly with either NPD or LPD
110 studs, with access *ad libitum* to the studs' respective diet. The presence of a vaginal plug the
following morning was taken as a sign of mating. Plug positive females were housed singly and
maintained on chow until offspring weaning at which time they were culled. At birth, offspring
were weighed and the litter male:female ratio determined. At 3 weeks of age all offspring were
weaned, the sexes caged separately per litter with access to chow and water *ad libitum* and allocated
115 randomly tail marks with permanent marker for subsequent weekly tracking of individuals. All
offspring were weighed weekly from birth till 24 weeks of age. All studs generated 2 litters each,
however, during pre-weaning development 2 NPD and 2 LPD litters, each from separate studs, were
lost due to maternal infanticide. As such, a total of 14 litters per dietary treatment were analysed.

120 ***Blood Pressure Measurement***

Blood pressure (systolic and diastolic) and heart rate were measured in stud males at 11
(pre-diet feeding), 17 (pre-mating) and 27 weeks of age, and all generated offspring from all 14
litters per treatment group (n = 42 NPD males, 43 NPD females, 27 LPD males and 42 LPD
females in total) at 6, 10, 14 and 18 weeks of age using a computerised, non-invasive, tail-cuff
125 system (Kent Scientific, USA). All mice were acclimatised to the experimental room for at least one
hour followed by a minimum of 30 minutes warming at 27-30°C prior to being placed within the
restraining and measurement apparatus for 5 minutes prior to measurement.

Glucose Tolerance Test (GTT)

130 Offspring glucose tolerance was determined in at least one male and female per litter at 22
weeks of age. Offspring were fasted overnight, with access to water *ad libitum*, and weighed
immediately prior to GTT. After administration of local anaesthetic (EMLA cream, Eutectic
Mixture of Local Anaesthetics, Lidocaine/Prilocaine, AstraZeneca, UK), fasting blood glucose
levels were determined in a sample collected from the tail vein using a hand-held glucometer

135 (Freestyle Optium, UK) prior to an intraperitoneal glucose bolus (2g/kg body weight in PBS).
Blood samples were collected from the tail vein at 15, 30, 60 and 90 minutes post-bolus for
determination of glucose concentration. All animals were returned to their original cage with
accesses to food and water *ad libitum*.

140 ***Mesenteric Artery Vasoreactivity***

Offspring (n = 8 pairs of male and female offspring per treatment group, each pair from
separate litters) vascular function was assessed at 24 weeks of age in isolated small mesenteric
artery segments as described previously (51) on a wire myograph (Danish Myo Technology A/S,
Denmark). Cumulative concentration response curves (CRCs) were measured for the α_1 -adrenergic
145 agonist phenylephrine (10^{-9} to 10^{-4} mol/L), and after submaximal (EC_{80}) pre-constriction with the
thromboxane mimetic U46619 (10 nmol/L), the vasodilators acetylcholine (ACh; 10^{-9} to 10^{-5}
mol/L) and isoprenaline (ISO; 10^{-10} to 10^{-6} mol/L) and the nitric oxide donor sodium nitroprusside
(SNP; 10^{-11} to 10^{-5} mol/L) in that order in the same arteries. All drugs were purchased from Sigma
(UK).

150

Tissue Sampling

All mice were culled by cervical dislocation. At 32 weeks of age, stud males were culled
and blood samples, taken via heart puncture, were allowed to clot on ice prior to centrifugation at
10,000 rpm, 4°C for 10 minutes, after which serum was aliquoted and stored at -80°C. Liver,
155 kidneys, heart, lungs, testes and retroperitoneal, gonadal, inguinal and interscapular fat (anatomical
locations defined previously (52)) were removed, weighed and stored at -80°C. Left and right
caudal epididymi were removed and placed within a pre-warmed 200 μ l drop of sperm motility
medium (135 mM NaCl, 5 mM KCl, 1 mM $MgSO_4$, 2 mM $CaCl_2$, 30mM HEPES; freshly
supplemented with 10 mM lactic acid, 1 mM sodium pyruvate, 20 mg/ml BSA, 25 mM $NaHCO_3$).
160 Epididymi were slashed several times using a 23 gauge needle and left for 15 minutes at 37°C for

sperm to swim out. A sample of sperm was taken for counting using a Neubaur counting chamber prior to assessment of motility. Collected sperm were pipetted under 2 ml of pre-warmed motility medium and left to swim up for one hour at 37°C. Sperm within the top 1.5 ml of medium were collected and counted as above. At 24 weeks of age, offspring were culled for collection of blood and somatic tissues (as described above).

Metabolite and Hormone Measurements

Following cull at 32 weeks of age, stud serum glucose was analysed using a commercial glucose oxidase assay (Sigma, UK) and serum insulin and testosterone levels determined by ELISA (Millipore and R&D Systems respectively, UK). Stud testes were homogenised (50 mM HEPES, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% SDS) prior to protein level determination (DC assay, Bio-Rad Laboratories, CA, USA). Testicular testosterone levels were determined using a commercial ELISA (R&D Systems). Levels of adiponectin and TNF- α were determined in offspring serum at cull by ELISA (R&D Systems). All assays were conducted in accordance with the manufacturer's instructions and measured on a Benchmark microplate reader (Bio-Rad Laboratories).

RNA extraction and transcript expression

RNA was extracted from offspring heart and liver tissues using the RNeasy Mini Kit (QIAGEN, UK) according to manufacturer's instructions. Contaminating genomic DNA was removed by on-column DNase I digestion prior to cDNA synthesis using the ImPromTMII kit (Promega, UK) using the included random primers. For Real-Time PCR (RTqPCR), 1 μ l (5 ng RNA equivalent) of cDNA was added to a mastermix comprising 10 μ l mastermix (2X Precision SYBRgreen Mastermix; PrimerDesign, UK), 0.7 μ l primer mix (5 μ M forward and reverse primers) and 8.3 μ l water per reaction. Water was used in place of cDNA as a no template control. Amplification and detection was performed using a Lightcycler 480 (Roche) and data acquired

using the LightCycler SW 1.5.lnk software. A post-amplification melting curve confirmed the presence of specific products for each primer set. Ct values were converted to relative expression values using the delta-delta Ct method with offspring heart data normalised to the expression of *Ppib* and *Sdha* and liver data normalised to *Pgk1* and *Tbp*. geNorm software (48) was used to determine these to be the most stable reference genes. Primer sequences and amplification efficiencies are provided in **Table 1**.

Statistical analyses

Where appropriate, stud male data were analysed using independent samples or repeated measures *t*-tests, following assessment for normality, and Pearson correlation for analysis of correlation between phenotypic measurements (SPSS version 17). Offspring litter sex ratios were analysed using a binomial test (SPSS version 17). Analysis of offspring vascular responsiveness was conducted using GraphPad Prism 6, with the log effective concentration equal to 50% of the maximal response (pEC50) and maximum response for each of the CRCs analysed with an independent samples *t*-test. All other offspring data were analysed using a multilevel random effects regression model (SPSS version 18) (53), accounting for paternal origin of litter, gestational litter size, offspring sex and body weight. Significance was taken at $P < 0.05$.

Results

Paternal LPD reduces stud growth

Prior to experimental feeding, there was no difference in mean NPD and LPD stud body weight (**Figure 1**). During the first 3 weeks of feeding, LPD stud males displayed a trend ($P < 0.1$) towards a lower body weight (i.e. at 12 -14 weeks of age), becoming lighter ($P < 0.05$) during weeks 4 - 9 (i.e. at 15 - 20 weeks of age, mating initiated at 18 weeks of age). Throughout the study, LPD studs grew more slowly ($P = 0.004$) (by 12%) than NPD studs. NPD and LPD stud systolic and diastolic blood pressure and heart rate did not differ throughout the entire study (data not shown).

Analysis of stud male organ weights at 32 weeks of age revealed a significantly lighter kidney (NPD 0.171 ± 0.004 g, LPD 0.149 ± 0.003 g; $P=0.0005$) and carcass (body weight minus the weight of collected organs and fat pads) (NPD 29.09 ± 0.27 g, LPD 27.59 ± 0.47 g; $P=0.015$) weight in LPD males. These differences remained when calculated as a proportion of body weight ($P < 0.05$; data not shown). No difference between NPD and LPD stud testosterone (combined serum and testicular; NPD 9.30 ± 5.24 ng/ml, LPD 11.40 ± 3.92 ng/ml) or serum insulin (NPD 2.26 ± 0.61 ng/ml, LPD 1.44 ± 0.12 ng/ml) were observed. However, serum glucose was higher in LPD studs (NPD 1.32 ± 0.08 mg/ml, LPD 1.55 ± 0.07 mg/ml) at a trend level ($P=0.06$). No differences in the total number of sperm (NPD $12.90 \times 10^6 \pm 2.30 \times 10^6$ /ml, LPD $11.62 \times 10^6 \pm 1.99 \times 10^6$ /ml), or number of sperm collected following swim-up (NPD $4.99 \times 10^6 \pm 1.29 \times 10^6$ /ml, LPD $4.68 \times 10^6 \pm 0.72 \times 10^6$ /ml) were observed between stud groups.

225 ***Paternal LPD affects offspring sex ratios, birth weight and adult phenotype***

Mean maternal weight prior to conception (16.28 ± 0.11 g), after 2 weeks of pregnancy (23.80 ± 0.29 g) and mean litter size (5.7 ± 0.4) did not differ between treatment groups. However, the proportion of male pups at birth was reduced in the LPD compared to the NPD treatment group (NPD 0.54 ± 0.04 , LPD 0.40 ± 0.06 ; $P=0.03$), and male offspring birth weight was increased (NPD 1.26 ± 0.02 g, LPD 1.33 ± 0.02 g; $P=0.05$). LPD offspring were also heavier than NPD offspring at 2 (NPD 6.54 ± 0.07 g, LPD 7.13 ± 0.08 g; $P=0.006$) and 3 (NPD 7.82 ± 0.10 g, LPD 8.61 ± 0.14 g; $P=0.019$) weeks of age. At weaning, the sexes were caged separately (mean of 3 NPD males, 2 LPD males, 3 NPD females and 3 LPD females per cage) with no further differences in body weights being observed between NPD and LPD offspring for up to 24 weeks of age (data not shown).

235 Analysis of correlations between paternal phenotype at the time of mating and offspring early postnatal development revealed significant negative correlations between LPD stud body weight and mean litter male:female ratio ($r=-0.444$, $P<0.0001$) and offspring weight at 1 ($r=-0.27$,

P =0.023) and 2 weeks of age ($r = -0.257$, $P = 0.030$). In NPD offspring, a positive correlation was
240 observed between stud body weight and litter male:female ratio ($r = 0.191$, $P = 0.085$) and offspring
weight at 2 weeks ($r = 0.210$, $P = 0.06$), which was significant at 1 week of age ($r = 0.224$, $P = 0.046$).
Additional negative correlations were observed between the number of days studs were on LPD and
litter male:female ratio ($r = -0.277$, $P = 0.018$), offspring birth weight ($r = -0.421$, $P < 0.0001$), weight
at 1 ($r = -0.363$, $P = 0.002$) and 2 weeks of age ($r = -0.348$, $P = 0.003$) which were not observed in
245 NPD offspring.

No difference in mean systolic or diastolic blood pressure or heart rates were observed
between NPD and LPD offspring at 6, 10 or 14 weeks of age (data not shown). However, at 18
weeks of age, LPD males displayed lower diastolic (NPD 84.80 ± 1.82 mmHg, LPD 75.88 ± 2.16
250 mmHg), systolic (NPD 113.13 ± 2.19 mmHg, LPD 103.93 ± 2.27 mmHg) and mean (NPD $93.93 \pm$
 1.91 mmHg, LPD 84.86 ± 2.16 mmHg) blood pressure, and elevated mean heart rate (NPD $682 \pm$
 11 beats per minute, LPD 711 ± 16 beats per minute; $P < 0.05$) (**Figure 2**).

At 22 weeks of age, both male and female LPD offspring displayed elevated blood glucose
255 concentrations following an intraperitoneal glucose bolus. At 15 and 60 minutes post injection, LPD
males had significantly elevated blood glucose concentrations, with a reduced overall clearance at
60 and 90 minutes (area under the curve, AUC; $P = 0.034$ and 0.029 respectively) (**Figure 3A**). LPD
females similarly had elevated blood glucose concentrations at 15 and 30 minutes post injection,
and impaired overall clearance at 60 and 90 minutes (AUC, $P = 0.022$ and 0.080 respectively)
260 (**Figure 3B**).

At 24 weeks of age, significantly attenuated vasoconstrictive responses (pEC50) to the α -1
adrenergic agonist phenylephrine (PE) and maximal vasodilatory responses to isoprenaline (ISO)
and the nitric oxide donor SNP were observed in arteries from LPD males (**Figure 4A**, $P < 0.05$).

265 Significantly attenuated pEC50 and maximal responses to SNP were also observed in LPD females
(**Figure 4B**, $P<0.05$).

Analysis of offspring organ and fat pad weights at 24 weeks of age revealed significantly increased inguinal (NPD 0.59 ± 0.04 g, LPD 0.77 ± 0.07 g; $P=0.017$) and total fat (combined
270 individual fat pads, NPD 2.02 ± 0.01 g, LPD 2.46 ± 0.18 g; $P=0.035$) weights in LPD males but not females. When expressed as a percentage of body weight, reduced heart (NPD 0.53 ± 0.01 , LPD 0.51 ± 0.02 ; $P=0.04$) and elevated inguinal fat (NPD 2.00 ± 0.12 , LPD 2.51 ± 0.20 ; $P=0.02$) and total fat (NPD 6.81 ± 0.34 , LPD 8.06 ± 0.53 ; $P=0.04$) proportions were observed in LPD males. As global adiposity levels influence metabolic state and glucose homeostasis, we performed additional
275 retrospective analyses incorporating body weight and adiposity measurements as ‘random effects’ within our regression analyses of offspring glucose tolerance. We observed a positive interaction between offspring body weight and overall glucose clearance (AUC) such that AUC increased by 40.57 per g increase in body weight ($P=0.004$), however, no interaction between total fat weight and AUC ($P=0.98$) was observed. In males, a significant positive interaction between body weight
280 and AUC was observed (AUC increased by 39.11 for each g increase in body weight; $P=0.04$), but no interaction with total fat and AUC ($P=0.41$) was present. Female offspring displayed no interaction between body weight and mean AUC ($P=0.48$), however, a positive interaction with total fat and AUC at a trend level (AUC increased by 219.76 for each g increase in total fat; $P=0.07$) was observed. No interaction between blood glucose and total adiposity were observed in
285 male or female offspring at each individual time point post glucose bolus. However, body weight was observed to interact positively with mean blood glucose levels in male offspring at 60 (0.621 mmol increase for every g increase in body weight; $P=0.017$) and 90 minutes (0.682 mmol increase for every g increase in body weight; $P=0.008$). No such interactions in female offspring were observed. Additional analyses of offspring adiposity and adult health revealed a negative correlation
290 in LPD offspring between BMI (weight g/length from nose to base of tail (cm^2)) and diastolic blood

pressure ($r = -0.276$, $P = 0.027$) and a positive correlation between total fat (g) and heart rate ($r = 0.279$, $P = 0.026$) at 18 weeks of age which were not present within NPD offspring. Serum adiponectin concentrations were greater ($P < 0.001$) in female than male offspring but were unaffected by paternal diet (**Table 2**). In contrast, there was a paternal-diet by offspring-sex interaction ($P = 0.015$) for TNF- α which indicated that this cytokine was elevated in LPD male compared to NPD male offspring, with no differences between paternal dietary treatments in female offspring. Analysis of correlation between adiponectin, TNF- α and offspring phenotype revealed significant negative correlations between adiponectin levels and body weight at cull in NPD and LPD offspring ($P < 0.0001$). Additionally, adiponectin levels correlated negatively with body weight at 3 weeks of age ($r = -0.384$; $P = 0.033$) and TNF- α levels ($r = -0.349$; $P = 0.05$) in LPD offspring. Finally, a positive correlation between TNF- α levels and body weight at cull ($r = 0.471$; $P = 0.008$) was also observed in LPD offspring.

Gene expression analyses in offspring heart for receptors involved in regulation of cardiac function revealed no differences for adrenergic receptor beta 1 (*Adrb1*), angiotensin II receptor type 1a (*Agtr1a*), bone morphogenetic protein receptor type II (*Bmpr2*), cholinergic receptor muscarinic 2 (*Chrm2*) or the solute carrier family 2 (facilitated glucose transporter), member 4 (*Glut4*) between NPD and LPD offspring (data not shown). However, analysis of genes involved in calcium signalling revealed significantly decreased expression of adenylate cyclase 5 (*Adcy5*; $P = 0.026$), phospholipase C beta1 (*Plcb1*; $P = 0.027$) and protein kinase C beta (*Prkcb*; $P = 0.008$) in offspring heart tissue (**Table 3**). As well as having a role in regulation of cardiovascular function, ADCY5, along with FTO (fat mass and obesity associated) have been identified as genes that are altered in type 2 diabetes (**3, 25**). Therefore, in response to the observations of impaired glucose tolerance and elevated adiposity in LPD offspring, we analysed the expression of *Adcy5* and *Fto* in offspring liver tissue. We observed no change in the expression of *Adcy5*, however, *Fto* was decreased ($P = 0.006$)

(**Table 3**). *Fto* expression in offspring cardiac tissue was also reduced ($P<0.001$) in NPD offspring (**Table 3**).

Discussion

320 To date, the majority of studies detailing the developmental programming of offspring health have focused on manipulation of the maternal environment. As such, our understanding of the impact of paternal nutrition on offspring development and long-term adult health remains poorly defined. In the present study, we have shown that LPD has minimal effects on paternal physiology and fertility (between 18 and 32 weeks of age), but that adult offspring derived from them display
325 significantly impaired cardiovascular and metabolic homeostasis. Our results provide evidence of an intergenerational modification of adult offspring phenotype in response to paternal diet.

Our results identify a series of offspring growth, cardiovascular and metabolic phenotypes whose regulation is compromised by paternal LPD. At birth, we observed a significantly reduced
330 litter male:female ratio and increased weight of male offspring from LPD studs. In mice, a low calorie diet fed to females results in a selective loss of male embryos during preimplantation development and subsequent skewing of litter sex ratios in favour of males (**38**). However, no such effects are observed when low calorie diets are fed to males. In contrast, no effects on litter sex ratio have been reported following maternal LPD in the mouse (**53, 54**). Based on previous reports (**38**),
335 we do not believe LPD induces a differential production in the number of X and Y bearing sperm in studs. However, differences in capacitation rates or motility between X and Y bearing sperm might explain these effects. Alternatively, LPD semen may induce a uterine environment more favourable to female preimplantation embryos, resulting in a selective loss of male embryos (**38**). However, additional studies are necessary to determine whether functional differences exist between X and Y
340 bearing sperm, and at what developmental stage offspring sex ratio is established.

Dysfunctional regulation between constriction and dilatation responses within resistance arteries has been identified in rodent models of cardiovascular programming (37, 51, 53). Augmentation of peripheral vascular function is characterised by the presence of altered endothelium-dependent vasodilation and/or changes in activity of signalling mechanisms regulating vascular smooth muscle function. We identified significant impairments in mesenteric artery responses to the α_1 -adrenergic agonist phenylephrine (PE), the β -adrenoreceptor agonist isoprenaline (ISO) and the nitric oxide donor sodium nitroprusside (SNP) in LPD offspring, however, no impairment in response to endothelial-dependent vasodilator acetylcholine (ACh) was observed. Within the resistance vasculature, endothelium-dependent vasodilatation is mediated predominantly through the action of endothelium-derived hyperpolarizing factor (EDHF) via small and intermediate calcium-activated potassium channels rather than NO (31). Indeed, in conditions of reduced NO bio-availability, up-regulation of EDHF activity has been observed (39). As ACh induces vascular smooth muscle cell hyperpolarization through both eNOS and EDHF pathways, a functional EDHF component would mask any impairment in NO signalling present. This could provide one mechanism through which altered responsiveness to SNP, but not ACh, could be manifest. Calcium homeostasis is central to the regulation of vascular smooth muscle function. PE induces vasoconstriction through the activation of phospholipase C and the mobilisation of calcium from intracellular stores, activating myosin light chain kinase. Conversely, isoprenaline elevates intracellular cAMP levels, activating protein kinase A, which inhibits myosin light chain kinase, causing vasodilatation. Isoprenaline also acts in concert with endothelial nitric oxide synthase, stimulating soluble guanylate cyclase to increase cGMP levels within the vascular smooth muscle, also inhibiting myosin light chain kinase activity. Therefore, impairment in vascular smooth muscle cell calcium signalling would impact negatively on both vaso-constriction and –dilatation function, as observed in LPD offspring. However, from the present study, the exact roles of NO, soluble guanylate cyclase, adenylate cyclase, prostaglandins and calcium-activated potassium channels are uncertain and would require further investigation.

Similarly, modulation of myocardial intracellular calcium signalling through altered
370 sympathetic or parasympathetic innervation, inhibition of calcium entry or AT1 receptor antagonists
(angiotensin II) would result in a lowering of blood pressure (56). As no difference in cardiac
expression of the β 1-adrenergic, angiotensin II type 1a or the cardiac muscarinic cholinergic
receptors were observed in offspring, we assessed the expression profiles of central regulators of
intracellular calcium signalling in offspring cardiac tissue. Here, we observed significantly
375 decreased expression of *Adcy5*, *Plcb*, and *Prkcb* in LPD offspring. Adenylate cyclase, in particular
Adcy5, plays a central role in modulating cardiac contractility, transducing the signal from the β -
adrenergic receptor, elevating cAMP-protein kinase A signalling and ultimately the influx of
calcium ions through voltage-dependent L-type calcium channels. Disruption of cardiac *Adcy5*
expression results in decreased in vivo responsiveness to isoprenaline, elevated heart rate through
380 reduced parasympathetic regulation, and reduced hypertrophy and apoptosis (29, 30). *Plcb* catalyses
the hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP2) to generate inositol (1,4,5)
triphosphate (IP3) and 1,2-diacylglycerol (DAG). IP3 initiates an increase in intracellular calcium,
whereas DAG activates protein kinase C beta (*Prkcb*). In turn *Prkcb*, a serine and threonine kinase,
phosphorylates a wide range of protein targets. Elevated expression of *Prkcb* isoforms, specifically
385 in the myocardium, is associated with hypertrophy, fibrosis, impairment of left ventricular
performance, progressive cardiomyopathy and heart failure (49, 55). In contrast, *Prkcb* inhibition
preserves cardiac contractility by attenuating diastolic dysfunction, myocyte hypertrophy, and
collagen deposition (12). Therefore, our observed relative hypotension, tachycardia and reduced
heart:body weight ratios observed in LPD offspring may result from impaired parasympathetic
390 stimulation of cardiac tissues providing decreased baroreflex restraint coupled with a reduced rate
of age related cardiomyopathy and fibrosis in response to reduced calcium signalling gene
expression (7), however, additional studies would be required to verify these conclusions.

In our current study, LPD offspring also exhibited reduced glucose tolerance and elevated adiposity in adulthood. Broad effects on the metabolism of young offspring following paternal nutritional manipulation have been reported. Offspring pancreatic β -cell function and gene expression are affected by paternal preconception high fat diet in rats (28), with paternal preconception fasting elevating offspring glucose levels in mice (2). Similarly, elevated expression of hepatic lipid and cholesterol biosynthesis genes, with decreased levels of cholesterol esters, at weaning have been observed in offspring mice from LPD fed studs (9), however, adult sex-specific cardiovascular, metabolic and glucogenic phenotype was not assessed. A causal link does exist between cardiovascular and metabolic phenotype, with increased adiposity and altered gene expression profiles, vascular dysfunction and hypertension being observed in female offspring from mouse dams fed a LPD exclusively during preimplantation development (52, 53). Impaired glycaemic homeostasis and increased adiposity are chronic inflammatory conditions associated with elevated levels of adipokines, inflammatory cytokines and oxidised low-density lipoproteins, all known to impair vascular smooth muscle and cardiac function (35, 46). Therefore, we measured the circulating levels of adiponectin and TNF- α in NPD and LPD offspring. Adiponectin levels correlate negatively with adiposity, with low levels being associated with cardio-metabolic disorders including endothelial dysfunction, type 2 diabetes and blood pressure (18). Conversely, elevated levels of the pro-inflammatory cytokine TNF- α are associated with insulin resistance and impact negatively on vascular function (4). We observed no significant difference in either adiponectin concentration between NPD and LPD males, or between NPD and LPD females. However, female offspring did display significantly higher adiponectin levels than males, reflective of previous reports (15). We also observed a significant paternal-diet by offspring-sex interaction indicating significant differences in the impact of paternal diet on offspring inflammatory responses dependent on sex. As such, the comparatively low levels of adiponectin, coupled with elevated levels of TNF- α , may contribute to the increased vascular dysfunction and impaired glucose homeostasis observed in LPD males. Conversely, in LPD females, the opposite relationship may

420 provide some protection against developing cardiovascular impairments to the same magnitude as those observed in the LPD males. Interestingly, we observed specific correlations between early postnatal body weight, adiponectin and TNF- α levels and adult body weight in LPD offspring. These data highlight the importance of early development and physiological characteristics (e.g. body weight) and adult markers of metabolic health, a central concept of the Developmental Origins of Health and Disease (DOHaD) hypothesis (21). Indeed, we have demonstrated previously similar 425 associations between early postnatal weight and adult cardiovascular dysfunction in a mouse maternal model of gestational LPD fed exclusively during preimplantation development (53). We also observed a negative correlation between BMI and diastolic blood pressure in LPD offspring, whilst total fat weight correlated positively with heart rate at 18 weeks of age, highlighting 430 additionally the interaction between adult adiposity and cardiovascular regulation within our model.

Human genome-wide associated studies of have identified significant associations between genetic polymorphisms and the risk prediction for type II diabetes including *ADCY5* and Fat Mass and Obesity associated gene (*FTO*) (1, 3). We observed significantly decreased expression of *Fto* in 435 the livers of LPD offspring, with no change in *Adcy5* expression. The *FTO* gene is an AlkB-like, Fe(II)- and 2-oxoglutarate-dependent nucleic acid demethylase, acting on single-stranded DNA and RNA, and has been showed to predispose individuals to diabetes through an effect on body mass index (43). In mice, fasting increases *Fto* expression in the liver within an inverse correlation between *Fto* mRNA and glucose levels (34), whilst knock out of the *Fto* gene results in reduced 440 lean mass and elevated fat mass (25). In mouse models of obesity, reduced hepatic *Fto* expression has been reported (44). As such, the reduced hepatic expression of *Fto* observed in NPD mice may disrupt regulation of energy metabolism, predisposing to impaired glucose tolerance and elevated adiposity.

445 Our observations raise the question as to the underlying developmental mechanisms through which paternal diet modifies adult offspring cardiovascular and metabolic phenotype. Recent studies have identified sperm hypomethylation and histone-enrichment at the promoters of developmental regulatory genes in both mice and men (8, 20). Carone et al., (9) also observed significant changes in the epigenome of sperm isolated from stud mice fed a LPD, correlating with weaning offspring metabolic phenotype. These observations highlight the potential that sperm epigenetic status could influence both sperm function and post fertilisation development and gene expression patterns. However, as the sperm epigenome is dramatically remodelled at fertilisation by the cytoplasm of the oocyte, the persistence of sperm epigenetic marks and their effects on offspring phenotype remains unknown. Recently, it has been demonstrated that paternal sensory environment prior to conception, influenced the sensory nervous system structure and function in F1 and F2 generations in mice (13). Interestingly, bisulfite sequencing analysis of paternal sperm and offspring tissue DNA revealed similar patterns of hypomethylation of the olfactory receptor *Olfir151*, proving potential evidence of an epigenetic basis of transgenerational inheritance of phenotype. Secondly, the relative contribution of sperm genomic-mediated programming and that determined by the composition of seminal plasma on long-term development and wellbeing of offspring remains to be established. It is known that seminal plasma cytokines stimulate maternal reproductive tract immunological responses, influencing embryonic, placental and offspring development (41). In human assisted reproductive cycles, there is an increasing awareness of the benefits of seminal plasma exposure on appropriate uterine responses and pregnancy outcomes following embryo transfer (47). Whether paternal LPD modifies the composition of the seminal plasma, and the impact this may have on uterine physiology following mating remains to be determined.

Our data extend the concept of developmental programming revealing the role of paternal nutrition in the early origins of adult offspring cardiovascular and metabolic health. These data are timely and relevant to human disease as paternal diet and lifestyle not only contribute to male-factor

infertility (5), but can also influence the cardiovascular and metabolic disease risk in subsequent generations (22, 32). Our observations highlight the need for a greater understanding of the underlying mechanisms through which parental diet and physiology affect gamete maturation, semen quality, and ultimately, long-term offspring health.

475

Acknowledgements

The authors wish to acknowledge Wing Yee Kwong for her technical assistance and advice.

This work was supported by a University of Nottingham Advanced Research Fellowship to Dr.

480 Adam Watkins and a Society for Reproduction and Fertility Academic Scholarship Fund award to Prof. Kevin Sinclair.

485 **Disclosures**

The authors declare no conflicts of interest, financial or otherwise.

References

1. **Abbas S, Raza ST, Ahmed F, Ahmad A, Rizvi S, and Mahdi F.** Association of Genetic polymorphism of PPARgamma-2, ACE, MTHFR, FABP-2 and FTO genes in risk prediction of type 2 diabetes mellitus. *J Biomed Sci* 20: 80, 2013.
2. **Anderson LM, Riffle L, Wilson R, Travlos GS, Lubomirski MS, and Alvord WG.** Preconceptional fasting of fathers alters serum glucose in offspring of mice. *Nutrition* 22: 327-331, 2006.
3. **Andersson EA, Pilgaard K, Pisinger C, Harder MN, Grarup N, Faerch K, Poulsen P, Witte DR, Jorgensen T, Vaag A, Hansen T, and Pedersen O.** Type 2 diabetes risk alleles near ADCY5, CDKAL1 and HHEX-IDE are associated with reduced birthweight. *Diabetologia* 53: 1908-1916, 2010.
4. **Aroor AR, McKarns S, Demarco VG, Jia G, and Sowers JR.** Maladaptive immune and inflammatory pathways lead to cardiovascular insulin resistance. *Metabolism* 62: 1543-1552, 2013.
5. **Bakos HW, Henshaw RC, Mitchell M, and Lane M.** Paternal body mass index is associated with decreased blastocyst development and reduced live birth rates following assisted reproductive technology. *Fertil Steril* 95: 1700-1704, 2011.
6. **Bellinger L, Lilley C, and Langley-Evans SC.** Prenatal exposure to a maternal low-protein diet programmes a preference for high-fat foods in the young adult rat. *Br J Nutr* 92: 513-520, 2004.
7. **Boyle AJ, Shih H, Hwang J, Ye J, Lee B, Zhang Y, Kwon D, Jun K, Zheng D, Sievers R, Angeli F, Yeghiazarians Y, and Lee R.** Cardiomyopathy of aging in the mammalian heart is characterized by myocardial hypertrophy, fibrosis and a predisposition towards cardiomyocyte apoptosis and autophagy. *Exp Gerontol* 46: 549-559, 2011.
8. **Brykczynska U, Hisano M, Erkek S, Ramos L, Oakeley EJ, Roloff TC, Beisel C, Schubeler D, Stadler MB, and Peters AH.** Repressive and active histone methylation mark distinct promoters in human and mouse spermatozoa. *Nat Struct Mol Biol* 17: 679-687, 2010.

9. **Carone BR, Fauquier L, Habib N, Shea JM, Hart CE, Li R, Bock C, Li C, Gu H, Zamore PD, Meissner A, Weng Z, Hofmann HA, Friedman N, and Rando OJ.** Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell* 143: 1084-1096, 2010.
10. **Ceelen M, van Weissenbruch MM, Prein J, Smit JJ, Vermeiden JP, Spreeuwenberg M, van Leeuwen FE, and Delemarre-van de Waal HA.** Growth during infancy and early childhood in relation to blood pressure and body fat measures at age 8-18 years of IVF children and spontaneously conceived controls born to subfertile parents. *Hum Reprod* 24: 2788-2795, 2009.
11. **Chavarro JE, Furtado J, Toth TL, Ford J, Keller M, Campos H, and Hauser R.** Trans-fatty acid levels in sperm are associated with sperm concentration among men from an infertility clinic. *Fertil Steril* 95: 1794-1797, 2011.
12. **Connelly KA, Kelly DJ, Zhang Y, Prior DL, Advani A, Cox AJ, Thai K, Krum H, and Gilbert RE.** Inhibition of protein kinase C-beta by ruboxistaurin preserves cardiac function and reduces extracellular matrix production in diabetic cardiomyopathy. *Circ Heart Fail* 2: 129-137, 2009.
13. **Dias BG, and Ressler KJ.** Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nat Neurosci* 2013.
14. **Edwards LJ, and McMillen IC.** Periconceptional nutrition programs development of the cardiovascular system in the fetal sheep. *Am J Physiol Regul Integr Comp Physiol* 283: R669-679, 2002.
15. **Eglit T, Lember M, Ringmets I, and Rajasalu T.** Gender differences in serum high-molecular-weight adiponectin levels in metabolic syndrome. *Eur J Endocrinol* 168: 385-391, 2013.
16. **Fernandez-Twinn DS, Wayman A, Ekizoglou S, Martin MS, Hales CN, and Ozanne SE.** Maternal protein restriction leads to hyperinsulinemia and reduced insulin-signaling protein expression in 21-mo-old female rat offspring. *Am J Physiol Regul Integr Comp Physiol* 288: R368-373, 2005.

17. **Ghanayem BI, Bai R, Kissling GE, Travlos G, and Hoffler U.** Diet-induced obesity in
540 male mice is associated with reduced fertility and potentiation of acrylamide-induced reproductive
toxicity. *Biol Reprod* 82: 96-104, 2010.
18. **Gu P, and Xu A.** Interplay between adipose tissue and blood vessels in obesity and vascular
dysfunction. *Rev Endocr Metab Disord* 14: 49-58, 2013.
19. **Hammoud AO, Gibson M, Stanford J, White G, Carrell DT, and Peterson M.** In vitro
545 fertilization availability and utilization in the United States: a study of demographic, social, and
economic factors. *Fertil Steril* 91: 1630-1635, 2009.
20. **Hammoud SS, Nix DA, Zhang H, Purwar J, Carrell DT, and Cairns BR.** Distinctive
chromatin in human sperm packages genes for embryo development. *Nature* 460: 473-478, 2009.
21. **Hanson M, Godfrey KM, Lillycrop KA, Burdge GC, and Gluckman PD.**
550 Developmental plasticity and developmental origins of non-communicable disease: theoretical
considerations and epigenetic mechanisms. *Prog Biophys Mol Biol* 106: 272-280, 2011.
22. **Kaati G, Bygren LO, and Edvinsson S.** Cardiovascular and diabetes mortality determined
by nutrition during parents' and grandparents' slow growth period. *Eur J Hum Genet* 10: 682-688,
2002.
- 555 23. **Kort HI, Massey JB, Elsner CW, Mitchell-Leef D, Shapiro DB, Witt MA, and
Roudebush WE.** Impact of body mass index values on sperm quantity and quality. *J Androl* 27:
450-452, 2006.
24. **Kwong WY, Wild AE, Roberts P, Willis AC, and Fleming TP.** Maternal undernutrition
during the preimplantation period of rat development causes blastocyst abnormalities and
560 programming of postnatal hypertension. *Development* 127: 4195-4202, 2000.
25. **McMurray F, Church CD, Larder R, Nicholson G, Wells S, Teboul L, Tung YC,
Rimington D, Bosch F, Jimenez V, Yeo GS, O'Rahilly S, Ashcroft FM, Coll AP, and Cox
RD.** Adult onset global loss of the *fto* gene alters body composition and metabolism in the mouse.
PLoS Genet 9: e1003166, 2013.

- 565 26. **Micke GC, Sullivan TM, Gatford KL, Owens JA, and Perry VE.** Nutrient intake in the bovine during early and mid-gestation causes sex-specific changes in progeny plasma IGF-I, liveweight, height and carcass traits. *Anim Reprod Sci* 121: 208-217, 2010.
27. **Mitchell M, Bakos HW, and Lane M.** Paternal diet-induced obesity impairs embryo development and implantation in the mouse. *Fertil Steril* 95: 1349-1353, 2011.
- 570 28. **Ng SF, Lin RC, Laybutt DR, Barres R, Owens JA, and Morris MJ.** Chronic high-fat diet in fathers programs beta-cell dysfunction in female rat offspring. *Nature* 467: 963-966, 2010.
29. **Okumura S, Kawabe J, Yatani A, Takagi G, Lee MC, Hong C, Liu J, Takagi I, Sadoshima J, Vatner DE, Vatner SF, and Ishikawa Y.** Type 5 adenylyl cyclase disruption alters not only sympathetic but also parasympathetic and calcium-mediated cardiac regulation. *Circ Res* 93: 364-371, 2003.
- 575 30. **Okumura S, Takagi G, Kawabe J, Yang G, Lee MC, Hong C, Liu J, Vatner DE, Sadoshima J, Vatner SF, and Ishikawa Y.** Disruption of type 5 adenylyl cyclase gene preserves cardiac function against pressure overload. *Proc Natl Acad Sci U S A* 100: 9986-9990, 2003.
31. **Ozkor MA, and Quyyumi AA.** Endothelium-derived hyperpolarizing factor and vascular function. *Cardiol Res Pract* 2011: 156146, 2011.
- 580 32. **Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K, Sjöström M, and Golding J.** Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 14: 159-166, 2006.
33. **Pisani LF, Antonini S, Pocar P, Ferrari S, Brevini TA, Rhind SM, and Gandolfi F.** Effects of pre-mating nutrition on mRNA levels of developmentally relevant genes in sheep oocytes and granulosa cells. *Reproduction* 136: 303-312, 2008.
- 585 34. **Poritsanos NJ, Lew PS, and Mizuno TM.** Relationship between blood glucose levels and hepatic Fto mRNA expression in mice. *Biochem Biophys Res Commun* 400: 713-717, 2010.
35. **Porter KE, and Riches K.** The vascular smooth muscle cell: a therapeutic target in Type 2 diabetes? *Clin Sci (Lond)* 125: 167-182, 2013.
- 590

36. **Rato L, Alves MG, Dias TR, Lopes G, Cavaco JE, Socorro S, and Oliveira PF.** High-energy diets may induce a pre-diabetic state altering testicular glycolytic metabolic profile and male reproductive parameters. *Andrology* 1: 495-504, 2013.
37. **Rodford JL, Torrens C, Siow RC, Mann GE, Hanson MA, and Clough GF.** Endothelial
595 dysfunction and reduced antioxidant protection in an animal model of the developmental origins of cardiovascular disease. *J Physiol* 586: 4709-4720, 2008.
38. **Rosenfeld CS.** Periconceptional influences on offspring sex ratio and placental responses. *Reprod Fertil Dev* 24: 45-58, 2011.
39. **Scotland RS, Madhani M, Chauhan S, Moncada S, Andresen J, Nilsson H, Hobbs AJ,
600 and Ahluwalia A.** Investigation of vascular responses in endothelial nitric oxide synthase/cyclooxygenase-1 double-knockout mice: key role for endothelium-derived hyperpolarizing factor in the regulation of blood pressure in vivo. *Circulation* 111: 796-803, 2005.
40. **Sinclair KD, Allegrucci C, Singh R, Gardner DS, Sebastian S, Bispham J, Thurston A, Huntley JF, Rees WD, Maloney CA, Lea RG, Craigon J, McEvoy TG, and Young LE.** DNA
605 methylation, insulin resistance, and blood pressure in offspring determined by maternal periconceptional B vitamin and methionine status. *Proc Natl Acad Sci U S A* 104: 19351-19356, 2007.
41. **Sjoblom C, Roberts CT, Wikland M, and Robertson SA.** Granulocyte-macrophage colony-stimulating factor alleviates adverse consequences of embryo culture on fetal growth
610 trajectory and placental morphogenesis. *Endocrinology* 146: 2142-2153, 2005.
42. **Soubry A, Schildkraut JM, Murtha A, Wang F, Huang Z, Bernal A, Kurtzberg J, Jirtle RL, Murphy SK, and Hoyo C.** Paternal obesity is associated with IGF2 hypomethylation in newborns: results from a Newborn Epigenetics Study (NEST) cohort. *BMC Med* 11: 29, 2013.
43. **Sovio U, Mook-Kanamori DO, Warrington NM, Lawrence R, Briollais L, Palmer CN,
615 Cecil J, Sandling JK, Syvanen AC, Kaakinen M, Beilin LJ, Millwood IY, Bennett AJ, Laitinen J, Pouta A, Molitor J, Davey Smith G, Ben-Shlomo Y, Jaddoe VW, Palmer LJ, Pennell CE,**

Cole TJ, McCarthy MI, Jarvelin MR, and Timpson NJ. Association between common variation at the FTO locus and changes in body mass index from infancy to late childhood: the complex nature of genetic association through growth and development. *PLoS Genet* 7: e1001307, 2011.

620 44. **Stratigopoulos G, Padilla SL, LeDuc CA, Watson E, Hattersley AT, McCarthy MI, Zeltser LM, Chung WK, and Leibel RL.** Regulation of Fto/Ftm gene expression in mice and humans. *Am J Physiol Regul Integr Comp Physiol* 294: R1185-1196, 2008.

45. **Todd SE, Oliver MH, Jaquiery AL, Bloomfield FH, and Harding JE.** Periconceptual undernutrition of ewes impairs glucose tolerance in their adult offspring. *Pediatr Res* 65: 409-413, 625 2009.

46. **Trayhurn P, and Wood IS.** Adipokines: inflammation and the pleiotropic role of white adipose tissue. *Br J Nutr* 92: 347-355, 2004.

47. **Tremellen KP, Valbuena D, Landeras J, Ballesteros A, Martinez J, Mendoza S, Norman RJ, Robertson SA, and Simon C.** The effect of intercourse on pregnancy rates during 630 assisted human reproduction. *Hum Reprod* 15: 2653-2658, 2000.

48. **Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, and Speleman F.** Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034, 2002.

49. **Wakasaki H, Koya D, Schoen FJ, Jirousek MR, Ways DK, Hoit BD, Walsh RA, and 635 King GL.** Targeted overexpression of protein kinase C beta2 isoform in myocardium causes cardiomyopathy. *Proc Natl Acad Sci U S A* 94: 9320-9325, 1997.

50. **Watkins AJ, and Fleming TP.** Blastocyst environment and its influence on offspring cardiovascular health: the heart of the matter. *J Anat* 215: 52-59, 2009.

51. **Watkins AJ, Lucas ES, Torrens C, Cleal JK, Green L, Osmond C, Eckert JJ, Gray 640 WP, Hanson MA, and Fleming TP.** Maternal low-protein diet during mouse pre-implantation development induces vascular dysfunction and altered renin-angiotensin-system homeostasis in the offspring. *Br J Nutr* 103: 1762-1770, 2010.

52. **Watkins AJ, Lucas ES, Wilkins A, Cagampang FR, and Fleming TP.** Maternal periconceptional and gestational low protein diet affects mouse offspring growth, cardiovascular and adipose phenotype at 1 year of age. *PLoS One* 6: e28745, 2011.

53. **Watkins AJ, Ursell E, Panton R, Papenbrock T, Hollis L, Cunningham C, Wilkins A, Perry VH, Sheth B, Kwong WY, Eckert JJ, Wild AE, Hanson MA, Osmond C, and Fleming TP.** Adaptive responses by mouse early embryos to maternal diet protect fetal growth but predispose to adult onset disease. *Biol Reprod* 78: 299-306, 2008.

54. **Watkins AJ, Wilkins A, Cunningham C, Perry VH, Seet MJ, Osmond C, Eckert JJ, Torrens C, Cagampang FR, Cleal J, Gray WP, Hanson MA, and Fleming TP.** Low protein diet fed exclusively during mouse oocyte maturation leads to behavioural and cardiovascular abnormalities in offspring. *J Physiol* 586: 2231-2244, 2008.

55. **Way KJ, Isshiki K, Suzuma K, Yokota T, Zvagelsky D, Schoen FJ, Sandusky GE, Pechous PA, Vlahos CJ, Wakasaki H, and King GL.** Expression of connective tissue growth factor is increased in injured myocardium associated with protein kinase C beta2 activation and diabetes. *Diabetes* 51: 2709-2718, 2002.

56. **Zubcevic J, Waki H, Raizada MK, and Paton JF.** Autonomic-immune-vascular interaction: an emerging concept for neurogenic hypertension. *Hypertension* 57: 1026-1033, 2011.

Figure Legends

Figure 1. Mean weekly body weight of NPD (open circles, n = 8) and LPD (closed circles, n = 8) stud males. Experimental feeding began at 11 weeks of age and continued till cull at 32 weeks. Mating to chow fed females began at 18 weeks of age. Error bars are S.E.M, *P<0.05.

Figure 2. Systolic (A), diastolic (B) and mean (C) blood pressure and heart rate (beats per minute; BPM) (D) in NPD (white bars) and LPD (black bars) offspring at 18 weeks of age. n = 34 NPD males, 36 NPD females, 24 LPD males and 42 LPD females. Error bars are S.E.M, *P<0.05.

Figure 3. Mean changes in blood glucose levels following an intraperitoneal glucose bolus (2g/kg body weight) in male (A), and female (B), NPD (open circles) and LPD (closed circles) offspring at 22 weeks of age. n = 15 male and 15 female offspring of each diet group representing all litters. Error bars are S.E.M, *P<0.05.

Figure 4. Mean vasoreactivity of isolated mesenteric arteries from male (A), and female (B), NPD (open circles) and LPD (closed circles) offspring at 24 weeks of age. Cumulative additions of phenylephrine (PE) and, after pre-constriction, of the vasodilators acetylcholine (ACh), isoprenaline (ISO) and sodium nitroprusside (SNP). n =8 males and 8 females of each diet group, each pair from separate litters. Error bars are S.E.M. *P<0.05.

Table 1. Real-time qPCR primer details.

Gene Name	Gene Symbol	Accession Number	<u>Primer Sequences</u>		Amplicon Length	Primer Efficiency
			Forward Primer	Reverse Primer		
Adrenergic receptor, beta 1	<i>Adrb1</i>	NM_007419.2	ggatgcctcttcgtcttct	cagtagatgatggggtgaagg	69	1.70
Angiotensin II receptor, type 1 alpha	<i>Agtr1a</i>	NM_177322.3	actcacagcaaccctccaag	ctcagacactgttcaaatgcac	62	1.84
Bone morphogenetic protein receptor, type II	<i>Bmpr2</i>	NM_007561.3	gagccctcccttgacctg	gtatcgaccccgccaatc	60	1.80
Cholinergic receptor, muscarinic 2, cardiac	<i>Chrm2</i>	NM_203491.3	tcggtgttaactgtcatcttc	tcaggttgaccggttg	109	1.85
Solute carrier family 2, member 4	<i>Glut4</i>	NM_009204.2	gacggacactccatctgttg	gccacgatggagacatagc	115	1.83
Adenylate cyclase 5	<i>Adcy5</i>	NM_001012765.4	atggaagctggtggcaag	cacctcatagccccattcag	78	1.70
Protein kinase C beta	<i>Prkcb</i>	NM_008855.2	aagcgagggcaatgaaga	cttctggagccttggtacctt	74	1.70
Phospholipase C beta 1	<i>Plcb1</i>	NM_001145830.1	tcgatgagaagcccaagc	ggcagcctttgaactgtc	67	1.75
Fat mass and obesity associated	<i>Fto</i>	NM_011936.2	tctgtctgccatcctggtc	tggtaaagccggacgactc	94	1.72
Phosphoglycerate kinase 1	<i>Pgk1</i>	NM_008828	tacctgctggctggatgg	cacagcctcgcatattct	65	1.70
Peptidylprolyl isomerase B	<i>Ppib</i>	NM_011149	ttctcataaccacagtcaagacc	acctccgtaccacatccat	92	1.80
Succinate dehydrogenase complex, subunit A, flavoprotein	<i>Sdha</i>	NM_023281	tgttcagttccacccaca	tctccacgacacccttctgt	66	1.88
TATA box binding protein	<i>Tbp</i>	NM_013684.3	gggagaatcatggaccagaa	gatgggaattccaggagtca	90	1.70

Table 2. Offspring serum adiponectin and TNF- α concentrations.

Sex	Males		Females		Significance (P)		
Diet	NPD	LPD	NPD	LPD	Diet	Sex	Diet x Sex
Adiponectin (ug/ml)	8.82 \pm 0.29	8.95 \pm 0.30	13.62 \pm 0.29	13.20 \pm 0.29	-	< 0.001	-
TNF- α (pg/ml)	4.02 \pm 1.82	9.38 \pm 1.89	5.63 \pm 1.86	3.64 \pm 1.86	-	-	0.015

Mean serum adiponectin and TNF- α concentrations. n = 16 males and 16 females from each dietary group, with all litters sampled. Values are mean \pm S.E.M.


690 **Table 3.** Offspring tissue transcript expression.

Tissue	Gene	NPD	LPD	P value
Heart	<i>Adcy5</i>	1.00 ± 0.02	0.93 ± 0.02	0.026
	<i>Fto</i>	1.00 ± 0.01	0.88 ± 0.02	<0.001
	<i>Plcb</i>	1.00 ± 0.03	0.89 ± 0.03	0.027
	<i>Prkcb</i>	1.00 ± 0.08	0.78 ± 0.03	0.008
Liver	<i>Adcy5</i>	1.00 ± 0.06	0.90 ± 0.06	0.285
	<i>Fto</i>	1.00 ± 0.02	0.90 ± 0.02	0.006

Mean relative transcript expression (± S.E.M.) for selected genes involved in calcium signalling and
695 metabolic regulation from NPD and LPD offspring heart and liver tissue. n = 10 males and 10
females from each dietary group, with each pair from separate litters. Transcript expression
normalised to that of *Ppib* and *Sdha* (heart) and *Pgk1* and *Tbp* (liver), and adjusted to NPD values
of 1.00.

700


Figure 1



705

710

Figure 2



715

720

725

730

Figure 3





Figure 4

